# C1498-eGFP-Puro



#### **Product Description**

Product Name: C1498-eGFP-Puro

Catalog Number: CL136 Lot Number: CL-IM141

Species: Mouse (Mus musculus)

Strain: C57BL/6J

Cell type: Lymphoblast (acute myeloid leukemia; AML)

Parental cells: C1498 (ATCC® TIB-49™)\*

Morphology: Lymphoblast Growth mode: Suspension

Reporter gene: Enhanced green fluorescent protein (eGFP)

Selection gene: Puromycin (Puro)

This is a cell line derived from the murine acute myeloid leukemia C1498 cell line (ATCC® TIB-49<sup>TM</sup>). Parental C1498 cells were transduced with LV-eGFP-PGK-Puro (Imanis #LV031) encoding the enhanced green fluorescent protein (eGFP) cDNA under the spleen focus-forming virus (SFFV) promoter and the puromycin resistance gene (Puro) under the phosphoglycerate kinase (PGK) promoter. A high eGFP expressing population was generated by selection using puromycin followed by selection using a methylcellulose based semi-solid medium. The lentiviral vector is a self-inactivating (SIN) vector in which the viral enhancer and promoter have been deleted. Transcription inactivation of the LTR in the SIN provirus increases biosafety by preventing mobilization by replication competent viruses and enables regulated expression of the genes from the internal promoters without *cis*-acting effects of the LTR<sup>1</sup>.

#### Mycoplasma Testing

This cell line has been tested for mycoplasma contamination and is certified mycoplasma free.

#### **Cell Line Authentication**

The parental C1498 cell line was authenticated and certified free of interspecies cross contamination by STR profiling.

#### **Recommended Uses**

These cells are suitable for *in vitro* and *in vivo* experimentation.

eGFP is not recommended for whole animal in-live imaging. Rather, samples can be collected post mortem for analysis by conventional fluorescence microscopy or flow cytometry.

#### References

<sup>1</sup>Miyoshi et al. J Virol. 1998. 72:8150-8157.

#### **Storage Instructions**

Remove cells from the dry ice packaging and immediately store in the vapor phase above liquid nitrogen (below -130°C).

#### **Complete Growth Medium**

High glucose DMEM 10% fetal bovine serum (FBS) 1% Penicillin/Streptomycin

Puromycin should <u>NOT</u> be added to the medium until a culture has been well established from the thawed cells (about 1 week). It is also recommended that a backup frozen cell stock be generated (see below) before adding puromycin to the growth medium.

Caution! Typical commercial puromycin stocks are provided at a concentration of 10 mg/mL or 10,000X.

To help maintain high eGFP expression, the cells can be subcultured in the presence of 1  $\mu$ g/mL puromycin. (Caution! Typical commercial puromycin stocks are provided at a concentration of 10 mg/mL or 10,000X.) Puromycin should not be added to the medium until a culture has been well established from the thawed cells (about 1 week). It is also recommended that a backup frozen cell stock be generated (see below) before adding puromycin to the growth medium.

### **Thawing Instructions**

- 1. Thaw cells by gently swirling in a 37°C water bath. To limit contamination, do not submerge the O-ring and cap.
- 2. When cells are ~70% thawed (less than 1 min), remove the vial and wipe down with 70% ethanol. Allow tube to dry completely.
- In a biosafety cabinet, transfer the cells into a 15 mL conical tube containing 5 mL of complete growth medium. Centrifuge cells at ~200 x g for 3-5 min.
- 4. Remove supernatant and resuspend cells in 1 mL complete growth medium. Remove an aliquot for counting.
- Dilute the cells further with growth medium to achieve a final density between 1 and 3 x 10<sup>6</sup> cells/mL (do NOT use more than 20 mL total). Transfer the cells to a T25 or T75 flask based on volume.
- 6. Incubate the culture at 37°C with 5% CO<sub>2</sub>.

### **Subculturing Instructions**

- Pipet the cell suspension gently to dislodge any cells loosely attached to the culture flask. Transfer the desired volume (half, one fifth, etc.) of the cells to a conical tube.
- 2. Centrifuge at ~150 x g for 3 min. (Note: a short, low speed spin is recommended to limit the amount of cell debris in the pellet.)
- 3. Remove supernatant and resuspend cells in complete growth medium. Transfer to an appropriate sized flask.

The cells should be subcultured as needed to maintain a density between 3 x  $10^5$  and 2 x  $10^6$  cells/mL. The cells can be passaged by dilution in fresh complete growth medium (without centrifugation). However, regular passage using centrifugation as described above is recommended to limit the amount of debris in cultures.

#### **Freezing Medium**

These cells can be amplified and used to generate additional frozen stocks. Preparation of low passage frozen stocks is highly recommended. Frozen stocks should be preserved in a designated cryopreservation medium or in complete growth medium without antibiotics supplemented with 5-10% DMSO.

<sup>\*</sup> The ATCC trademark and any and all ATCC catalog numbers are trademarks of the American Type Culture Collection

# C1498-eGFP-Puro




#### **Certificate of Analysis**

Testing performed by Imanis Life Sciences

Test description	Result
Post thaw viable cell recovery	84%
Cells per vial	~ 2 x 10 <sup>7</sup>
Sterility	No contamination detected
Mycoplasma	No contamination detected
Puromycin selection	Pass QC
Fluorescence expression	Pass QC
Average doubling time	26.6 h*


<sup>\*</sup>Doubling time represents the average doubling time during logarithmic growth. This value should be used for general estimation only.

#### Morphology



Cell photos taken at 200x magnification.

#### Fluorescence Expression



C1498-eGFP-Puro (green) or isotype control (C1498-Fluc-Puro; grey) cells were fixed with paraformaldehyde and analyzed by flow cytometry.

## **Legal Disclaimers**

LIMITED PRODUCT WARRANTY
THIS WARRANTY LIMITS OUR LIABILITY TO REPLACEMENT OF THIS PRODUCT. NO OTHER WARRANTES OF ANY KIND, EXPRESS OR
IMPLIED, INCLUDING, WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE
PROVIDED BY IMANIS. IMANIS SHALL HAVE NO LIABILITY FOR ANY DIRECT, INDIRECT, CONSEQUENTIAL, OR INCIDENTAL DAMAGES
ARISING OUT OF THE USE, THE RESULTS OF USE, OR THE INABILITY TO USE THIS PRODUCT.

FOR IN VITRO USE ONLY. THIS CERTIFICATE IS A DECLARATION OF ANALYSIS AT THE TIME OF MANUFACTURE.

#### PURCHASER NOTIFICATION

#### LIMITED LICENSE NOTICE - RESEARCH USE ONLY

THE IMANIS CELL LINES ARE NOT INTENDED FOR USE IN HUMANS. CELL LINES TRANSDUCED WITH LENTIVIRAL VECTORS ARE CLASSIFIED AS BIOSAFETY LEVEL 2 REAGENTS AND SHOULD BE USED UNDER THE APPROPRIATE BIOSAFETY LEVEL PER INSTITUTIONAL GUIDELINES.

THE PURCHASER AGREES THAT IMANIS MATERIALS DESIGNATED AS BIO-SAFETY LEVEL 2 CONSTITUTE KNOWN PATHOGENS AND THAT OTHER IMANIS MATERIALS NOT SO DESIGNATED AND ANY PROGENY OR MODIFICATION MAY BE PATHOGENIC UNDER CERTAIN CONDITIONS. PURCHASER ASSUMES ALL RISK AND RESPONSIBILITY IN CONNECTION WITH THE RECEIPT, HANDLING, STORAGE, DISPOSAL, TRANSFER AND USE OF THE IMANIS MATERIALS INCLUDING WITHOUT LIMITATION TAKING ALL APPROPRIATE SAFETY AND HANDLING PRECAUTIONS TO MINIMIZE HEALT HOR ENVIRONMENTAL RISK PURCHASER AGREES THAT ANY ACTIVITY UNDERTAKEN WITH THE IMANIS MATERIALS AND ANY PROGENY OR MODIFICATION WILL BE CONDUCTED IN COMPLIANCE WITH ALL APPLICABLE GUIDELINES, LAWS AND REGULATIONS.

THE IMANIS MATERIAL, ANY OTHER IMANIS PRODUCTS, AND ANY TECHNICAL INFORMATION AND ASSISTANCE PROVIDED BY IMANIS ARE PROVIDED "AS IS", WITHOUT WARRANTIES OF ANY IMPLE SEARCH STATE OF MERCHANDED TO ANY IMPLIED WARRANTIES OF MERCHANTED TO ANY IMPLIED WARRANTIES OF MERCHANTED TO ANY IMPLIED TO

IN NO EVENT SHALL MANIS, ITS PARENTS, SUBSIDIARIES, DIRECTORS, OFFICERS, AGENTS, EMPLOYEES, ASSIGNS, SUCCESSORS AND AFFILIATE (COLLECTIVELY "IMANIS INDEMNIFIED PARTIES") BE LIABLE FOR INDIRECT, SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES OF ANY KIND IN CONNECTION WITH OR ARISINS OUT OF THIS AGREEMENT (WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, STATUTE OR OTHERWISE) EVEN IF IMANIS HAS BEEN ADVISED, KNEW OR SHOULD HAVE KNOWN OF THE POSSIBILITY OF SUCH DAMAGES, INCLUDING, BUT NOT LIMITED TO, LOST PROFITS, COST OF CAPITAL, COST OF SUBSTITUTE PRODUCTS OR CLAIMS OF LUCENSEES CUSTOMERS FOR SUCH DAMAGE. IN NO EVENT SHALL IMANIS CUMULATIVE LIABILITY EXCEED THE ACTUAL AMOUNTS OF LICENSEES AND ADDRESS THIS CREENEST OF SHALL AMOUNTS OF THE CONTRACT OF T

Quality control by: AMW **Quality Assurance by: RLV** Effective Date: 26-Oct-2022