Nalm6-Fluc-Puro/HuCD19-Puro (Low)



Product Description

Product Name: Nalm6-Fluc-Puro/HuCD19-Puro (Low)

Catalog Number: CL182 Lot Number: IMP034

Species: Human (*Homo sapiens*)
Tissues: Peripheral blood

Cell type: Lymphoma

Parental cells: Nalm6 (ATCC® CRL-3273™)*

Morphology: Lymphocyte-like Growth mode: Suspension

Reporter gene: Firefly luciferase (Fluc)

Transgene: CD19

Selection gene: Puromycin (Puro)

This is a cell line derived from the human B cell precursor leukemia Nalm6 cell line (ATCC® CRL-3273TM). Imanis Nalm6-Fluc-Puro cells (Imanis #CL151) were transduced with a lentiviral vector encoding the CD19 cDNA and the puromycin resistance gene. A low CD19 expressing population was generated by selection using a semi-solid methyl-cellulose-based medium. The lentiviral vectors are self-inactivating (SIN) vectors in which the viral enhancer and promoter have been deleted. Transcription inactivation of the LTR in the SIN provirus increases biosafety by preventing mobilization by replication competent viruses and enables regulated expression of the genes from the internal promoters without *cis*-acting effects of the LTR (Miyoshi et al. J Virol. 1998. 72:8150-8157).

* The ATCC trademark and any and all ATCC catalog numbers are trademarks of the American Type Culture Collection

Mycoplasma Testing

This cell line has been tested for mycoplasma contamination and is mycoplasma free.

Cell Line Authentication

Authentication of the parental Nalm6 cell line was confirmed by STR profiling.

Recommended Uses

These cells are suitable for *in vitro* and *in vivo* experimentation.

The Fluc transgene facilitates *in vivo* noninvasive bioluminescent imaging of implanted cells.

References

¹Miyoshi et al. J Virol. 1998. 72:8150-8157.

Biosafety Notice

This cell line was generated by transduction with a lentiviral vector. Cell lines transduced with lentiviral vectors are classified as biosafety level 2 reagents and should be used under appropriate biosafety level for institutional guidelines.

Storage Instructions

Remove cells from the dry ice packaging and immediately store in the vapor phase above liquid nitrogen (below -130°C).

Complete Growth Medium

RPMI-1640 Medium (RPMI) supplemented with 10 mM HEPES 10% Fetal Bovine Serum (FBS) 1% Penicillin/Streptomycin 1 µg/mL Puromycin

Puromycin should <u>NOT</u> be added to the medium until a culture has been well established from the thawed cells (about 1 week). It is also recommended that a backup frozen cell stock be generated (see below) before adding puromycin to the growth medium.

Caution! Typical commercial puromycin stocks are provided at a concentration of 10 mg/mL or 10,000X.

Thawing Instructions

- 1. Thaw cells by gently swirling in a 37°C water bath. To limit contamination, do not submerge the O-ring and cap.
- 2. When cells are ~70% thawed (less than 1 min), remove the vial and wipe down with 70% ethanol. Allow tube to dry completely.
- 3. In a biosafety cabinet, transfer the cells into a 15 mL conical tube containing 5 mL of complete growth medium. Centrifuge cells at ~300 x g for 3-5 min.
- Remove supernatant and resuspend cells in 1 mL complete growth medium. Remove an aliquot for counting.
- Dilute the cells further with growth medium to achieve a final density of 1 x 10⁶ cells/mL. Transfer the cells to a T25 or T75 flask based on volume.
- 6. Incubate the culture at 37°C with 5% CO₂. Cells should reach full confluency 3 days after thawing.
 - Note: For optimal cell health and viability, change media every ~ 3 days even if the cells aren't confluent.

Subculturing Instructions

The cells should be subcultured through dilution in fresh complete growth medium as needed to maintain a density between 7×10^5 and 3×10^6 cells/mL. As needed, passage using centrifugation as described below to limit the amount of debris in cultures.

- 1. Pipet the cell suspension gently to dislodge any cells loosely attached to the culture flask. Transfer the desired volume (half, one fifth, etc.) of the cells to a conical tube.
- 2. Centrifuge at ~150 x g for 3 min. (Note: a short, low speed spin is recommended to limit the amount of cell debris in the pellet.)
- 3. Remove supernatant and resuspend cells in complete growth medium. Transfer to an appropriately sized flask.

Freezing Medium

These cells can be amplified and used to generate additional frozen stocks. Preparation of low passage frozen stocks is highly recommended. Frozen stocks should be preserved in a designated cryopreservation medium or in complete growth medium without puromycin supplemented with 5-10% DMSO.

Nalm6-Fluc-Puro/HuCD19-Puro (Low)



Certificate of Analysis

Testing performed by Imanis Life Sciences


Test description	Result
Post thaw viable cell recovery	83%
Cells per vial	~ 8.5 x 10 ⁶
Sterility	No contamination detected
Mycoplasma	No contamination detected
Puromycin selection	Pass QC
Luciferase expression	Pass QC
Human CD19 expression	Pass QC
Average doubling time	40.7 h*

^{*}Doubling time represents the average doubling time during <u>logarithmic growth</u>. This value should be used for general estimation only.

High density, 200X


Morphology






Low- and high-density photos taken at different times after thawing

Luciferase Expression




The indicated number of cells were placed in wells of a 96-well plate. After the addition of 15 mg/mL d-luciferin, bioluminescence was immediately read using a microplate reader.


Human CD19 Expression





Nalm6-Fluc-Puro/HuCD19-Puro (Low) cells were stained with an anti-HuCD19 antibody (blue) or isotype control antibody (grey) and analyzed by flow cytometry.

Expression Profiling of Surface Markers









Nalm6-Fluc-Puro/HuCD19-Puro (Low) (blue) cells were stained with an anti-HuBCMA antibody (A), anti-HuCD20 antibody (B), anti-HuCD38 antibody (C), anti-HuGPRC5D antibody (D), or isotype control antibody (grey; all panels) and analyzed by flow cytometry.

Quality control by: AWD Quality Assurance by: RLV Effective Date: 13-Nov-2024

Legal Disclaimers

THE IMANIS CELL LINES ARE NOT INTENDED FOR USE IN HUMANS. CELL LINES TRANSDUCED WITH LENTIVIRAL VECTORS ARE CLASSIFIED AS BIOSAFETY LEVEL 2 REAGENTS AND SHOULD BE USED UNDER THE APPROPRIATE BIOSAFETY LEVEL PER INSTITUTIONAL GUIDELINES.

LIMITED PRODUCT WARRANT

THIS WARRANTY LIMITS OUR LIABILITY TO REPLACEMENT OF THIS PRODUCT. NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING, WITHOUT LIMITATION, IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE PROVIDED BY IMANIS. IMANIS SHALL HAVE NO LIABILITY FOR ANY DIRECT, INDIRECT, CONSEQUENTIAL, OR INCIDENTAL DAMAGES ARISING OUT OF THE USE, THE RESULTS OF USE, OR THE IMBILITY TO USE THIS PRODUCT.

FOR IN VITRO USE ONLY. THIS CERTIFICATE IS A DECLARATION OF ANALYSIS AT THE TIME OF MANUFACTURE.

PURCHASER NOTIFICATION

LIMITED LICENSE NOTICE - RESEARCH USE ONLY

THE PURCHASER AGREES THAT IMANIS MATERIALS DESIGNATED AS BIO-SAFETY LEVEL 2 CONSTITUTE KNOWN PATHOGENS AND THAT OTHER IMANIS MATERIALS NOT 50 DESIGNATED AND ANY PROGENY OR MODIFICATION MAY BE PATHOGENIC UNDER CERTAIN CONDITIONS, PURCHASER A SSUMES ALL RISK AND RESPONSIBILITY IN CONNECTION WITH THE RECEIPT, HANDING, STORAGE, DISPOSAL, TRANSFER AND USE OF THE IMANIS MATERIALS INCLUDING WITHOUT LIMITATION TAKING ALL APPROPRIATE SAFETY AND HANDLING PRECAUTIONS TO MINIMIZE HEALTH OF ENVIRONMENTAL RISK, PURCHASER AGREES THAT ANY ACTIVITY UNDERTRAKEN WITH THE IMANIS MATERIALS AND ANY PROGENY OR MODIFICATION WILL BE CONDUCTED IN COMPLIANCE WITH ALL APPLICABLE GUIDELINES, LAWS, AND REQUILATIONS.

THE IMANIS MATERIAL, ANY OTHER IMANIS PRODUCTS, AND ANY TECHNICAL INFORMATION AND ASSISTANCE PROVIDED BY IMANIS ARE PROVIDED AS IS, WITHOUT WARRANTIES OF ANY KIND, EXPRESS OR IMPLED, INCLUDING BUT NOT LIMITED TO ANY IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, MANUFACTURE ACCORDING TO 6MP STANDARDS, TYPICALITY, SAFETY, ACCURACY AND NON-INFRINGEMENT.

IN NO EVENT SHALL IMANIS, ITS PARENTS, SUBSIDIARIES, DIRECTORS, OFFICERS, AGENTS, EMPLOYEES, ASSIGNS, SUCCESSORS AND AFFILLATE COLLECTIVELY "MANIS INDEMNIFIED PARTIES") BE LIABLE FOR INDIRECT, SPECIAL, INCIDENTAL ON ENSEQUENTIAL DAMAGES OF ANY KIND IN CONNECTION WITH OR ARISING OUT OF THIS AGREEMENT (WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, STATUTE OR OTHERWISE) EVEN IF IMANIS HAS BEEN ADVISED, KNEW OR SHOULD HAVE KNOWN OF THE POSSIBILITY OF SUCH DAMAGES, INCLUDING, BUT NOT LIMITED TO, LOST PROFITS, COST OF CAPITAL, COST OF SUBSTITUTE PRODUCTS OR CLAIMS OF LICENSEE'S CUSTOMERS FOR SUCH DAMAGE. IN NO EVENT SHALL IMANIS CUMULATIVE LIABILITY EXCEED THE ACTUAL AMOUNTS PAID BY PURCHASER UNDER THIS AGREEMENT FOR THE TWENT (12) MONTH PERIOD PRECEDING THE DATE OF THE EVENT GIVING RISE TO THE CLAIM. THE PROVISIONS OF THIS SECTION SHALL SURVIVE THE EXPIRATION OR TERMINATION OF THIS AGREEMENT AND SHALL APPLY EVEN IF THE LIMITED REMEDS PRECIFIED IN THIS AGREEMENT IS AGREEMENT IS AGREEMENT IN SAGREMENT IS SOUND TO HAVE FALLED OF ITS ESSENTIAL PURPOSE.